

nVH Italia srl[®]

Herbal extracts manufacturers



BroccOX[®]

BRASSICA OLERACEA DRY EXTRACT 0.3% - 0.5% SULFORAPHANE

BROCCOX® - Brassica oleracea dry extract 0.3%-0.5% sulforaphane

Origin of the drug: Italy
Origin of the extract: Italy
Solvent: Water
Full traceability: from beginning to end

DOSAGE:

Food grade extract: 200-250 mg/die
Cosmetic grade extract: 0,5-2%

Food grade extract:

- **Indirect antioxidant:** broccoli is itself rich in antioxidants such as carotenoids, flavonoids, vitamin C, isothiocyanates. Due to the presence of cysteine and methionine, it also encourages the production of glutathione, an important endogenous antioxidant
- **Induction of detoxification:** an important property of the Brassica family is the capacity to "switch on" genes that produce cellular enzymes required for optimum cell function. These enzymes convert toxins into water-soluble substances that are readily excreted.
- **Antibacterial and antifungal :** sulforaphane and other glucosinolates have a powerful inhibitory effect on *Helicobacter pylori*, a bacterium found in the stomach of many individuals.

Cosmetic grade extract :

- **Protection skin from sun:** a cream made from young broccoli shoots protects the skin against the harmful effects of UV radiation. Sulforaphane incites skin cells to produce enzymes that protect the skin from within. The symptoms of burns turn out to be high on 40% less pronounced in people who have applied cream containing broccoli. Furthermore, this protection remains in effect for several days
- **Seboregulation and skin conditioner :** its content in mineral salts is important for maintaining skin hydrated and for regulation sebum

BROCCOX® is 100% Made in Italy from beginning (RAW MATERIAL) to end (DRY EXTRACT)

BIBLIOGRAPHY

1. Nestle M. Broccoli sprouts as inducers of carcinogen-detoxifying enzyme systems: clinical, dietary, and policy implications. *Proc Natl Acad Sci U S A*. 1997;94(21):11149-51.
2. 2-Minich DM. Blending science of the clinical efficacy and safety of cruciferous vegetable phytochemicals. *Nutr Rev*. 2007;65(6 Pt 1):259-67.
3. 3- Weng JR, Tai CH, Kuo SM, et al. Indole-3-carbinol as a chemopreventive and anti-cancer agent. *Cancer Lett*. 2008;262(2):153-63.
4. 4- Kack AS, Finley JW. Cruciferous vegetables: cancer protective mechanisms of glucosinolate hydrolysis products and selenium. *Integr Cancer Ther*. 2004;3(1):5-12.
5. 5- Matusheshi NV, Jeremy EH. Comparison of the bioactivity of two glucoraphanin hydrolysis products found in broccoli, sulforaphane and sulforaphane nitrile. *J Agric Food Chem*. 2001;49(12):5743-9.
6. 6- Rouzaud C, Yousef SA, Duncan AJ. Hydrolysis of glucosinolates to isothiocyanates after ingestion of raw or microwaved cabbage by human volunteers. *Cancer Epidemiol Biomarkers Prev*. 2004;13(11):25-31.
7. 7- Sharma A, Sharma NK, Madhupratibha SV, et al. Targeting Akt3 signaling in malignant melanoma using isoselenocyanates. *Clin Cancer Res*. 2009;15(9):1674-85.
8. 8- Nayan-Ashraf MH, Sadeghinejad Z, Davies GF, et al. Phase 2 protein inducers in the diet promote healthier aging. *J Gerontol A Biol Sci Med Sci*. 2008;63(11):1168-76.
9. 9- Fahey JW, Zhang Y, Talalay P. Broccoli sprouts: an exceptionally rich source of inducers of enzymes that protect against chemical carcinogens. *Proc Natl Acad Sci U S A*. 1997;94(42):10367-72.
10. 10- Brooks JD, Paton VG, Vidanes G. Potent induction of phase 2 enzymes in human prostate cells by sulforaphane. *Cancer Epidemiol Biomarkers Prev*. 2001;10(9):949-54.
11. 11- Dinkova-Kostova AT, Talalay P. Direct and indirect antioxidant properties of inducers of cytoprotective proteins. *Mol Nutr Food Res*. 2008;52 Suppl 1:S128-38.
12. 12- Heits E, Herhaus C, Klimo K, et al. Nuclear factor kappa B is a molecular target for sulforaphane-mediated anti-inflammatory mechanisms. *J Biol Chem*. 2001;276(54):2008-15.
13. 13- Moon DO, Kim MO, Kang SH, et al. Sulforaphane suppresses TNF-alpha-mediated activation of NF-kappaB and induces apoptosis through activation of reactive oxygen species-dependent caspase-3. *Cancer Lett*. 2009;274(1):132-42.
14. 14- Nayan-Ashraf MH, Sadeghinejad Z, Juurlink BH. Dietary approach to decrease aging-related CNS inflammation. *Nutr Neurosci*. 2005;8(2):101-10.
15. 15- Haristoy X, Fahey JW, Scholtus L, et al. Evaluation of the antimicrobial effects of several isothiocyanates on *Helicobacter pylori*. *Planta Med*. 2005;71(9):336-30.
16. 16- Cheney G. Vitamin U therapy of peptic ulcer. *Calif Med*. 1952;77(4):248-52.
17. 17- Johansson NL, Pavia CS, Chiao JW. Growth inhibition of a spectrum of bacterial and fungal pathogens by sulforaphane, an isothiocyanate product found in broccoli and other cruciferous vegetables. *Planta Med*. 2008;74(7):747-50.
18. 18- Thejas P, Kuttan G. Immunomodulatory activity of Sulforaphane, a naturally occurring isothiocyanate from broccoli (Brassica oleracea). *Phytochemistry*. 2007;14(7-8):538-45.
19. 19- Thejas P, Kuttan G. Modulation of cell-mediated immune response in B16-F1 melanoma-induced metastatic tumor-bearing C57BL/6 mice by sulforaphane. *Immunopharmacol Immunotoxicol*. 2007;29(2):173-86.
20. 20- Michnovicz JJ, Adlercreutz H, Bradlow HL. Changes in levels of urinary estrogen metabolites after oral indole-3-carbinol treatment in humans. *J Natl Cancer Inst*. 1997;89(10):718-23.

nVH Italia srl[®]

Your partner from beginning to end

Via Cellini, 16/A - 22071 - Cadorago - Co - Italy
Phone +39 031 905071 | Fax +39 031 904869
email info@nvhextracts.com
web www.nvhextracts.com

