

nVH Italia srl

Herbal extracts manufacturers

FOOD & COSMETIC



CapsiRED®

CAPSICUM FRUIT $\geq 2.0\%$ CAPSAICINOIDS EXPRESSED AS CAPSAICIN
BEADLETS

CAPSIRE® - Capsicum fruit >= 2.0% capsaicinoids expressed as capsaicin BEADLETS

Origin of the drug: Italy
Origin of the extract: Italy
Solvent: Water/ Ethanol

Full traceability: from beginning to end

DOSAGE:

Food grade extract: 200 mg/die

Cosmetic grade extract: 0.025%- 0.3% antiinflammatory

Food grade extract:

- **Thermogenesis activity** : thermogenesis is the process by which the body produces heat. The adaptive thermogenesis is controlled by the sympathetic nervous system through which certain hormones (especially norepinephrine) produced by the adrenal glands and muscle, inducing increased production of heat. Capsaicin is able to increase the availability of norepinephrine and therefore the increase of lipolysis for the production of heat
- **Prevention and healing of gastric ulcers**: Capsaicin inhibits acid secretion, stimulates alkali and mucus secretion and particularly gastric mucosal blood flow which helps to prevent gastric ulcers.
- **Hypocholesterolemic**: prevention of cholesterol gallstones and protection of the structural integrity of erythrocytes under conditions of hypercholesterolemia.

Cosmetic grade extract:

- **Reversive and vasodilating**: topically, Capsaicin has the ability to reactivate the microcirculation and blood flow to the tissues. It's useful in the formulation of products for the treatment of joint and muscle pain, in preparations for the care of scalp to stimulate their growth and even to the revitalizing effect in products for skin.

CAPSIRE® is 100% Made in Italy from beginning (RAW MATERIAL) to end (BEADLETS)

BIBLIOGRAPHY

- Segal KR, Gutin B. "thermic effects of food and exercise in lean and obese women". Metabolism 1983; 32:581 Dulloo et al. International Journal of Obesity 2001; 522-529
- Rutledge J., Toxicological review of Capsaicinoids (PDF), 1993
- V.S. Govindarajan, M.N. Sathyanarayana, Capsicum - Production, Technology, Chemistry and Quality, Part I - V, 1985 - 1991
- Frenzuoli F. Fitoroterapia. IV ed. Milano: Elsevier, 2008. 3
- Appendino G, De Petrocellis L, Trevisani M, et al. Development of the first ultra-potent "capsaicinoid" agonist at transient receptor potential vanilloid type 1 (TRPV1) channels and its therapeutic potential. J Pharmacol Exp Ther 2005; 312: 561-70.
- Backonja M, Wallace MS, Blonsky ER, et al. NGX-4010, a high-concentration capsaicin patch, for the treatment of postherpetic neuralgia: a randomised, double-blind study. Lancet Neurol 2008; 7: 1106-12.
- Mason L, Moore RA, Derry S, Edwards JE, McQuay HJ. Systematic review of topical capsaicin for the treatment of chronic pain. BMJ 2004; 328: 991.
- Ellison N, Loprinzi CL, Kugler J, et al. Phase III placebo-controlled trial of capsaicin cream in the management of surgical neuropathic pain in cancer patients. J Clin Oncol 1997; 15: 2974-80.
- Epstein JB, Marcocci JH. Topical application of capsaicin for treatment of oral neuropathic pain and trigeminal neuralgia. Oral Surg Oral Med Oral Pathol 1994; 77: 135-40.
- Low PA, Dpfel-Gehring TL, Dyck PJ, Litchy WJ, O'Brien PC. Double-blind, placebo-controlled study of the application of capsaicin cream in chronic distal painful polyneuropathy. Pain 1995; 62: 163-8.
- De Sio M, Gallien P, Denys P, et al. Intravesical glucidic capsaicin versus glucidic solvent in neurogenic detrusor overactivity: a double blind controlled randomized study. NeuroUrol Urodyn 2006; 25: 752-7.
- Lazzari M, Benfiori P, Benigni G, Maggi CA, Lecci A, Turini D. Intravesical capsaicin for treatment of severe bladder pain: a randomized placebo controlled study. J Urol 1996; 156: 947-52.
- Bortolotti M, Coccia G, Grossi G, Miglioli M. The treatment of functional dyspepsia with red pepper. Aliment Pharmacol Ther 2002; 16: 1075-82.
- Geppetti P, Nassini R, Materazzi S, Benemei S. The concept of neurogenic inflammation. BJU Int 2008; 101 (Suppl 3): 2-4.
- Marabini S, Ciabatti PG, Poli G, Fusco BM, Geppetti P. Beneficial effects of intranasal applications of capsaicin in patients with vasomotor rhinitis. Eur Arch Otorhinolaryngol 1991; 248: 191-4.
- Ciabatti PG, D'Acunzio L. Intranasal Capsicum spray in idiopathic rhinitis: a randomized prospective application regimen trial. Acta Otolaryngol 2008; 14: 1-5.
- Frenzuoli F. Interazioni tra erbe, alimenti e farmaci. II ed. Milano: Tecniche Nuove, 2009. 1
- Tuntipopparat S, Judprasong K, Zeder C, et al. Chili, but not turmeric, inhibits iron absorption in young women from an iron-fortified composite meal. J Nutr 2006; 136: 2970-4.

nV Italia srl
Your partner from beginning to end

Via Cellini, 16/A - 22071 - Cadorago - Co - Italy
Phone +39 031 905071 | Fax +39 031 904869
email info@nvhextracts.com
web www.nvhextracts.com

