

Sherri S. Levin, M.D. & Associates

Sherri S. Levin, M.D. Amelie Chu, M.D.

Sooyoung Hwang, M.D. Katherine Whitty-Schnitzer, M.D.

OBSTETRICS • GYNECOLOGY • INFERTILITY

Phone: 713-464-4111 • Fax 713-464-3116

Congratulations! Your expected due date is _____.

Patient portal:

Please be sure to register on our patient portal. You will find the link in your email that you have provided to us. Once you have signed in, you will receive all future appointment confirmations and all normal lab results. We will no longer be calling with appointment reminders or **normal lab results**. We are encouraging all our patients to use the patient portal to contact our office instead of calling and leaving a message. You will be able to request prescriptions, schedule appointments and send secure emails through the patient portal. We will be able to respond to your portal message faster than with a phone call. We ask you **NOT** to leave urgent messages on the portal. You will not receive any unsolicited or marketing emails from us.

Approximate timeline of events for your pregnancy

16 weeks: Screening for Down's Syndrome or Spina bifida (lab draw)

3D ultrasound may be scheduled for fun!

20 weeks: Anatomy Ultrasound with technician in our office.

Your doctor will discuss the results immediately afterwards with you.

24 weeks: 1-hour glucose test instructions given.

28 weeks: 1-hour glucose test should be completed.

Information will be given on how to register with the hospital. This process may be started online, but must be completed in person on the 3rd floor. Also, information on classes offered will be given, if you are interested.

36 weeks: Screening for GBS (culture) HIV & syphilis (hospital mandated lab draw)

Lab work:

_____ **NATERA LABS: These labs should be scheduled on or after _____**

1. Panorama non-invasive prenatal test- Chromosome conditions tested: Trisomy 21, Trisomy 18, Trisomy 13 and fetal sex if desired)
2. Horizon Carrier Screening- Checks to see if you are a carrier of one of four autosomal recessive or X-Linked genetic conditions. (Cystic Fibrosis, Duchenne Muscular Dystrophy, Fragile X Syndrome and Spinal Muscular Atrophy) This testing is recommended by The American Congress of Obstetricians and Gynecologist. (ACOG)

Recommended Vaccines in pregnancy:

Tdap / Whooping Cough Vaccine (Between 27 weeks and before delivery.)

Flu Vaccine

Rhogam at 28 weeks if your blood type is a negative.

Welcome to Dr. Sherri Levin and Associates! We are very excited to care for you during your pregnancy and we want you to have a healthy pregnancy. This packet reviews some basic conditions and information about pregnancy but it is *not* comprehensive and you are welcome to read additional pregnancy guides that are helpful to you.

Your pregnancy should be shared with the family and you may bring other children to your appointments if you desire. Small children should have another adult to supervise them during your appointment when you have your ultrasound (around 20 weeks).

Our clinic hours are as follows:

Monday 8:00-4:00

Tuesday 8:30-4:00

Wednesday 8:00-4:00

Thursday 8:00-4:00

Friday 8:00-12:00

Closed on Saturday and Sunday and some major holidays.

If you call the office when it is closed, you will be given the number for the answering service. If you have concerns that you feel need to be discussed with a doctor, please call and leave a message with the answering service. Your phone call will be returned by one of practices physicians; Dr. Levin, Dr. Gonzalez, Dr. Chu, or Dr. Hwang depending on who is on call.

General prenatal instructions

During your pregnancy you should abstain from alcohol; medications or drugs that are not safe in pregnancy or that have not been cleared by your doctor, smoking and douching or use of tampons. You should limit your caffeine intake to 200 mg daily (approximately 12 oz coffee).

Common medical problems

***Heartburn** – You may use TUMS, Maalox, or Mylanta. If you are taking more than the recommended amount or feel like your symptoms are not improving, you may take Zantac (ranitidine) over the counter as recommended on the bottle. If your symptoms do not improve, you should talk to your doctor about other available treatments.

***Colds**- You may take over-the-counter medications (listed in the chart) for symptom relief. You should call the office if your temperature goes over 101 F or if your symptoms worsen. You should call the office if you have asthma or other medical problems that make you more susceptible to breathing difficulties.

***Headaches** - Acetaminophen (Tylenol) is safe to take in pregnancy (as recommended on the bottle). If you have a severe headache that is sudden in onset and not relieved with acetaminophen you should call the office.

***Constipation-** Constipation is a common problem during pregnancy because the hormones slow down the movement in the GI system. It is very important to drink 6-8 glasses of water daily and to increase your fiber intake to approximately 25 gm per day. You can eat fiber-rich foods or take a fiber supplement such as Metamucil, Citrucel, Fibercon and Benefiber. It is also safe to take Colace (docusate) which is a stool softener. Occasional use of Milk of Magnesia is safe.

***Hemorrhoids-** Can also develop or worsen during pregnancy due to the increased pressure on the veins in the rectum from the pregnant uterus. It is important not to strain during a bowel movement. Warm baths and Preparation H or other similar hemorrhoid cream can help reduce symptoms.

Nausea and Vomiting of Pregnancy (Morning sickness)

Nausea often begins in the early weeks of pregnancy and worsens by weeks 10-12 and gradually improves after that for most women. It can occur at any time during the day. It is important to eat frequent small meals. Bland carbohydrates are well tolerated as well as citrus juices. Occasional vomiting is not uncommon but you should call the office if you are unable to keep any fluids down for a 24-hour period. Please see the medication list for medications that can be taken over-the-counter to treat symptoms.

Dental problems

Dental problems can occur and should be taken care of when they arise. X-rays are allowed given you are shielded properly with a lead shield. We can provide you a dental authorization upon request.

Swelling

Swelling can occur at any time during pregnancy but worsens in the third trimester. It is important to stay well hydrated and to avoid a high sodium diet. It also helps to elevate the legs and feet above the level of the heart as much as possible while at home. Ted hose are tight compression stockings which also help reduce swelling.

Exercise in pregnancy and the postpartum period

If you do not have any medical or obstetric complications, 30 minutes or more of moderate exercise a day on most, if not all, days of the week is recommended (150 minutes per week). After the first trimester you should avoid lying on your back or standing motionless for long periods of time. You may need to be monitored more closely if you engage in very strenuous physical activity during pregnancy (such as long-distance running). Contact sports or activities that have an increased risk of falling should be avoided. Scuba diving should be avoided. The changes that occur in your body during pregnancy generally persist for 4-6 weeks postpartum and you should wait until after this period to return to your pre-pregnancy exercise routine or until cleared by your doctor.

Travel

Unless you have been told by your doctor that it is not safe to travel, occasional air travel during pregnancy is considered safe and is allowable up to 28 weeks. If you are traveling via air you may also need documentation of your gestational age which we can provide for you.

Zika Virus: (Zika) is a disease caused by Zika virus that is spread to people primarily through the bite of an infected *Aedes* species mosquito. The most common symptoms of Zika are fever, rash, joint pain, and conjunctivitis (red eyes). The illness is usually mild with symptoms lasting for several days to a week after being bitten by an infected mosquito. People usually don't get sick enough to go to the hospital, and they very rarely die of Zika. For this reason, many people might not realize they have been infected. Once a person has been infected, he or she is likely to be protected from future infections. The CDC suggests DEET products to prevent insect bites.

What is DEET?

DEET (chemical name, N-diethyl-meta-toluamide) is used as the active ingredient in many insect repellents. Insect repellents that contain DEET offer the best protection against mosquito bites. DEET is designed for direct application to skin to repel insects, rather than kill them. Products containing DEET currently are available to the public in a variety of liquids, lotions, sprays, and impregnated materials (e.g., wrist bands). Concentrations of DEET in products that are designed for application to skin range from 4% to 100%.

What does the DEET concentration mean?

The concentration of DEET in a product indicates how long the product will be effective. A higher concentration does not mean that the product will work better; it means that it will be effective for a longer period of time. Therefore, products containing lower concentrations of DEET may need to be reapplied, depending on the length of time a person is outdoors.

Please refer to the CDC web site listed below, for the most current locations that should be avoided. <http://www.cdc.gov/zika/geo/active-countries.html>

***Please understand that complications can occur during any pregnancy and if you have a complication while traveling in the US or out of the country, it may be unsafe for you to travel home. If you deliver prematurely, it may be a prolonged period of time before it is safe for you or the baby to return home. For these reasons, we recommend staying in the local area in the third trimester.**

When you travel, it is important to minimize the risk of a venous thromboembolism (blood clot) and you should move your lower extremities or get up and walk around every two hours (as long as it is safe). Compression stockings should also be worn to reduce swelling while traveling; as well as to reduce the risk of a blood clot. If you have been on a long trip and you notice pain and swelling in one leg you should call the office immediately.

Diet and Weight Gain

The recommended weight gain in pregnancy is different for each individual based on a woman's pre-pregnancy body mass index (BMI). We will review this with you at one of your first appointments and give you a sheet outlining your target weight gain. If you have any of the following: BMI >30, family history of diabetes in a first degree relative, history of gestational diabetes in a previous pregnancy or if you have had a previously large baby (over 4000 gm/8 lb 8oz). If you do not have any of the above, we will do a routine glucose test around 28 weeks to screen for gestational diabetes. We will recommend an early glucose screen to insure you do not have diabetes or insulin resistance.

Sexual activity

You may continue to engage in sexual activity as long as it remains comfortable for you, and you have not been told to abstain by your doctor. If you have any of the following: cerclage in place, preterm labor or ruptured membranes. There may also be other conditions in which sexual activity should be avoided and your doctor will discuss this with you. It is normal to contract after sexual activity or an orgasm but they should go away with rest. The cervix is very soft in pregnancy; It is not uncommon to have some spotting after intercourse but you should call the office if it is bright red bleeding. You should avoid all sexual activity and avoid putting anything in the vagina

Symptoms of Miscarriage (First trimester)

If you have active bleeding, passage of tissue or clots or severe cramping accompanied by bleeding you should call the office. If you pass tissue you should try to place it in a container so that it can be brought into the office. It is not always necessary to be seen in the emergency room for a miscarriage and the doctor on call will decide the appropriate treatment for you. If you have severe pain; light headiness, and have not yet had an ultrasound to confirm a normal pregnancy in the uterus you should call the office immediately. *It is important to understand that 1/3 of all pregnancies have some bleeding in the first trimester and this does not necessarily mean you are having a miscarriage.*

Concerning Symptoms in Second and Third Trimester

Call the office immediately or go to Labor and Delivery for the following: gush of fluid or bright red bleeding (like a period).

It is normal to have Braxton-Hicks contractions in the second and third trimester. They are usually irregular. If they become persistent you should rest, drink lots of water to ensure that you are well hydrated and ensure that you are emptying your bladder regularly. Start timing them from the start of one contraction to the start of another contraction. If you have 8 contractions in an hour you should call the office. You may find that a maternity/belly band may be beneficial.

If you have other symptoms such as abdominal pain, worsening headaches, changes in your vision or any other concerning symptoms, please call the office.

Cord blood banking

It is the law in Texas to inform you of umbilical cord blood banking. Umbilical cord blood contains cells that can be transplanted into a person to aid in the treatment of some diseases. There are public and private cord blood banks. There is rigorous screening for infectious diseases prior to donation to a public bank. Private banks were initially designed for use by the same individual later in life, should that individual develop a disease. The chance of this unit of blood being used by the individual or family member is approximately 1/2700.

Foods to Avoid in Pregnancy

All poultry, meat and fish should be cooked thoroughly. Avoid seafood high in mercury such as swordfish, shark, king mackerel and tilefish. It is safe to eat 8-12 ounces of seafood every week such as; shrimp, crab, canned light tuna (chunk white tuna and tuna steak should be limited to 6 ounces per week), salmon, pollock, catfish, cod and tilapia. Avoid all raw fish and raw shellfish. Cook eggs thoroughly. Avoid unpasteurized foods. Many cheeses are very healthy and safe to eat in pregnancy such as mozzarella and cottage cheese. Cheeses such as brie, feta, camembert, blue cheese and Mexican-style cheeses such as queso fresco are not always pasteurized and should be avoided. If the cheese is labeled as pasteurized or made with pasteurized milk, it is safe in pregnancy. All deli meats that are kept in the refrigerator (and do not need to be heated) and that are not expired are also fine to eat.

Additional information provided for you today should include:

- ❖ Drugs that are safe in pregnancy
- ❖ Immunizations and Pregnancy

Safe Medications

During pregnancy, women can be more susceptible to ailments like cold and flu and other conditions. Only certain medications are safe during pregnancy. The following are considered relatively safe, but you should use these very sparingly, especially decongestants of any kind. Prescription medications should be taken exactly as directed and you should check with us before starting any new prescription. Follow the labels for dosage and directions.

ACNE Benzoyl Peroxide Clindamycin Topical Erythromycin Salicylic Acid (Avoid in 3rd trimester) AVOID: Accutane Retin-A Tetracycline Minocycline	ANTIBIOTICS Ceclor Cephalosporins E-mycin Keflex Macrobid/ Macroclantin Penicillin Zithromax AVOID: Cipro Tetracycline Minocycline Levaquin Bactrim	COLDS/ALLERGIES Benadryl, Claritin, Zyrtec Chlor-Trimeton, Dimetapp Mucinex(guaifenesin) Sudafed**/Sudafed-12 Hour** Tylenol Cold & Sinus** Vicks Vapor Rub **AVOID if problems with blood pressure
CONSTIPATION MiraLAX, Colace, Dulcolax Suppository, Metamucil, Milk of Magnesia	COUGH Cough Drops Robitussin (Plain & DM)	LICE RID AVOID: Kweil
GAS Gas-X Mylicon Phazyme	HEADACHES Cold Compress Tylenol/ Acetaminophen Caffeine (up to 200mg per day)	HEARTBURN Avoid lying down for at least 1 hour after meals Aciphex, Maalox, Mylanta, Pepcid, Prilosec, Rolaids, Zantac, Tums, Omeprazole, Lansoprazole
HEMORRHOIDS Hydrocortisone, OTC Preparation H Tucks pads	HERPES Acyclovir Valtrex/ Valacyclovir	NASAL SPRAY Saline Nasal Spray Flonase
NAUSEA Vitamin B6 25mg 3 times daily Unisom ¼ or ½ tablet at bedtime Dramamine Emetrol Ginger Root 250mg 4 times a day Diclegis	PAIN Tylenol	RASH Benadryl Hydrocortisone Cream
SLEEP AIDS Benadryl, Chamomile Tea, Unisom, Tylenol PM	THROAT Cepacol Cepastat Throat Lozenges	TOOTH PAIN Oragel Can be treated by Dentist
YEAST INFECTION Monistat-3 Terazol-3		