


WHAT TO EXPECT AFTER TREATMENT



- For the majority of patients, downtime with **BELKYRA™** is minimal and occurs most commonly after their first treatment session.¹
- Apply ice or a cold pack to the treatment area for 10 to 15 minutes as needed.¹
- The most common side effects are swelling, bruising, pain, numbness, redness, areas of hardness, and tingling or itchiness in the treatment area. You can find detailed Important Safety Information on the reverse side.²
- While unlikely, talk to your doctor if you develop an asymmetric smile, facial muscle weakness, difficulty swallowing, or if symptoms worsen.²
- Remember, once your goals are achieved, retreatment with **BELKYRA™** is not expected.¹


belkyra™
deoxycholic acid injection



PUT YOUR **BEST CHIN** FORWARD

Actual results with BELKYRA™.

Many people treated with **BELKYRA™** experienced visible improvement in the appearance of their chin profile in just 2 to 4 treatments.^{*2}

Up to 6 treatments may be administered.²

*Spaced at least one month apart. ¹Unretouched patient photos taken before and after 3 treatment sessions with **BELKYRA™**. Patient did not experience significant weight loss or gain between the start and end of treatment. Individual results will vary.¹



My Next Appointment

Date: _____ Time: _____

Before your next treatment, ask your doctor if there are any special instructions you should follow to prepare for your appointment.