



# What to expect FROM TREATMENT



*b*  
Pr **belkya**<sup>™</sup>  
deoxycholic acid injection

- ✓ For most patients, downtime with BELKYRA<sup>™</sup> was minimal and occurred most commonly after the first treatment session<sup>1</sup>
- ✓ The most common side effects are swelling, bruising, pain, numbness, redness, and areas of hardness, tingling or itchiness in the treatment area. These are not all the side effects of BELKYRA<sup>™</sup>. Please see Important Safety Information on the other side. Talk to your doctor if you develop an asymmetric smile, facial muscle weakness, difficulty swallowing, or if any existing symptom worsens.<sup>2</sup>
- ✓ Apply ice or a cold pack to the treatment area for 10 to 15 minutes as needed<sup>1</sup>
- ✓ Talk to your doctor if you have questions or concerns about managing side effects<sup>2</sup>

## ACTUAL RESULTS WITH BELKYRA<sup>™</sup>



MANY PATIENTS TREATED  
WITH BELKYRA<sup>™</sup> EXPERIENCED  
VISIBLE CONTOURING OF  
THEIR CHIN PROFILE IN  
**2 TO 4 TREATMENTS.**<sup>†</sup>  
UP TO 6 TREATMENTS  
MAY BE ADMINISTERED.<sup>2</sup>

Unretouched photos of clinical trial patient taken before and after 4 treatment sessions with BELKYRA<sup>™</sup>.

Sex: Female Age: 35 Weight (before)\*: 115.5 lbs Weight (after)\*: 113.3 lbs Total Dose: 16.2 mL Individual results may vary.<sup>1</sup>

\*Patient did not experience significant weight loss or gain between the start and end of treatment.

<sup>†</sup>Spaced at least one month apart.

## My next

### APPOINTMENT

DATE \_\_\_\_\_ DAY \_\_\_\_\_ TIME \_\_\_\_\_

PRIOR TO YOUR NEXT TREATMENT, ASK YOUR DOCTOR IF THERE ARE ANY SPECIAL INSTRUCTIONS YOU SHOULD FOLLOW TO PREPARE FOR THE APPOINTMENT.